

First Observation of $\Upsilon(1S) \rightarrow \gamma\pi\pi$

CLEO Collaboration
(February 7, 2008)

Abstract

We report on a study of exclusive radiative decays of the $\Upsilon(1S)$ resonance collected with the CLEO II detector operating at CESR. We present the first observation of the radiative decays $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ and $\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0$. For the dipion mass regime $m_{\pi\pi} > 1.0$ GeV we obtain $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-) = (6.3 \pm 1.2 \pm 1.3) \times 10^{-5}$ and $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0) = (1.7 \pm 0.6 \pm 0.3) \times 10^{-5}$. The observed $\gamma\pi\pi$ events are consistent with the hypothesis $\Upsilon(1S) \rightarrow \gamma f_2(1270)$.

A. Anastassov,¹ J. E. Duboscq,¹ K. K. Gan,¹ T. Hart,¹ K. Honscheid,¹ H. Kagan,¹ R. Kass,¹ J. Lee,¹ H. Schwarthoff,¹ A. Wolf,¹ M. M. Zoeller,¹ S. J. Richichi,² H. Severini,² P. Skubic,² A. Undrus,² M. Bishai,³ S. Chen,³ J. Fast,³ J. W. Hinson,³ N. Menon,³ D. H. Miller,³ E. I. Shibata,³ I. P. J. Shipsey,³ S. Glenn,⁴ Y. Kwon,⁴ Permanent address: Yonsei University, Seoul 120-749, Korea. A.L. Lyon,⁴ S. Roberts,⁴ E. H. Thorndike,⁴ C. P. Jessop,⁵ K. Lingel,⁵ H. Marsiske,⁵ M. L. Perl,⁵ V. Savinov,⁵ D. Ugolini,⁵ X. Zhou,⁵ T. E. Coan,⁶ V. Fadeyev,⁶ I. Korolkov,⁶ Y. Maravin,⁶ I. Narsky,⁶ R. Stroynowski,⁶ J. Ye,⁶ T. Wlodek,⁶ M. Artuso,⁷ E. Dambasuren,⁷ S. Kopp,⁷ G. C. Moneti,⁷ R. Mountain,⁷ S. Schuh,⁷ T. Skwarnicki,⁷ S. Stone,⁷ A. Titov,⁷ G. Viehhauser,⁷ J.C. Wang,⁷ J. Bartelt,⁸ S. E. Csorna,⁸ K. W. McLean,⁸ S. Marka,⁸ Z. Xu,⁸ R. Godang,⁹ K. Kinoshita,⁹ I. C. Lai,⁹ P. Pomianowski,⁹ S. Schrenk,⁹ G. Bonvicini,¹⁰ D. Cinabro,¹⁰ R. Greene,¹⁰ L. P. Perera,¹⁰ G. J. Zhou,¹⁰ S. Chan,¹¹ G. Eigen,¹¹ E. Lipeles,¹¹ J. S. Miller,¹¹ M. Schmidtler,¹¹ A. Shapiro,¹¹ W. M. Sun,¹¹ J. Urheim,¹¹ A. J. Weinstein,¹¹ F. Würthwein,¹¹ D. E. Jaffe,¹² G. Masek,¹² H. P. Paar,¹² E. M. Potter,¹² S. Prell,¹² V. Sharma,¹² D. M. Asner,¹³ J. Gronberg,¹³ T. S. Hill,¹³ D. J. Lange,¹³ R. J. Morrison,¹³ H. N. Nelson,¹³ T. K. Nelson,¹³ D. Roberts,¹³ B. H. Behrens,¹⁴ W. T. Ford,¹⁴ A. Gritsan,¹⁴ H. Krieg,¹⁴ J. Roy,¹⁴ J. G. Smith,¹⁴ J. P. Alexander,¹⁵ R. Baker,¹⁵ C. Bebek,¹⁵ B. E. Berger,¹⁵ K. Berkelman,¹⁵ V. Boisvert,¹⁵ D. G. Cassel,¹⁵ D. S. Crowcroft,¹⁵ M. Dickson,¹⁵ S. von Dombrowski,¹⁵ P. S. Drell,¹⁵ K. M. Ecklund,¹⁵ R. Ehrlich,¹⁵ A. D. Foland,¹⁵ P. Gaidarev,¹⁵ R. S. Galik,¹⁵ L. Gibbons,¹⁵ B. Gittelman,¹⁵ S. W. Gray,¹⁵ D. L. Hartill,¹⁵ B. K. Heltsley,¹⁵ P. I. Hopman,¹⁵ J. Kandaswamy,¹⁵ D. L. Kreinick,¹⁵ T. Lee,¹⁵ Y. Liu,¹⁵ N. B. Mistry,¹⁵ C. R. Ng,¹⁵ E. Nordberg,¹⁵ M. Ogg,¹⁵ Permanent address: University of Texas, Austin TX 78712. J. R. Patterson,¹⁵ D. Peterson,¹⁵ D. Riley,¹⁵ A. Soffer,¹⁵ B. Valant-Spaight,¹⁵ A. Warburton,¹⁵ C. Ward,¹⁵ M. Athanas,¹⁶ P. Avery,¹⁶ C. D. Jones,¹⁶ M. Lohner,¹⁶ C. Prescott,¹⁶ A. I. Rubiera,¹⁶ J. Yelton,¹⁶ J. Zheng,¹⁶ G. Brandenburg,¹⁷ R. A. Briere,¹⁷ A. Ershov,¹⁷ Y. S. Gao,¹⁷ D. Y.-J. Kim,¹⁷ R. Wilson,¹⁷ H. Yamamoto,¹⁷ T. E. Browder,¹⁸ Y. Li,¹⁸ J. L. Rodriguez,¹⁸ S. K. Sahu,¹⁸ T. Bergfeld,¹⁹ B. I. Eisenstein,¹⁹ J. Ernst,¹⁹ G. E. Gladding,¹⁹ G. D. Gollin,¹⁹ R. M. Hans,¹⁹ E. Johnson,¹⁹ I. Karliner,¹⁹ M. A. Marsh,¹⁹ M. Palmer,¹⁹ M. Selen,¹⁹ J. J. Thaler,¹⁹ K. W. Edwards,²⁰ A. Bellerive,²¹ R. Janicek,²¹ P. M. Patel,²¹ A. J. Sadoff,²² R. Ammar,²³ P. Baringer,²³ A. Bean,²³ D. Besson,²³ D. Coppage,²³ C. Darling,²³ R. Davis,²³ S. Kotov,²³ I. Kravchenko,²³ N. Kwak,²³ L. Zhou,²³ S. Anderson,²⁴ Y. Kubota,²⁴ S. J. Lee,²⁴ R. Mahapatra,²⁴ J. J. O'Neill,²⁴ R. Poling,²⁴ T. Riehle,²⁴ A. Smith,²⁴ M. S. Alam,²⁵ S. B. Athar,²⁵ Z. Ling,²⁵ A. H. Mahmood,²⁵ S. Timm,²⁵ and F. Wappler²⁵

¹Ohio State University, Columbus, Ohio 43210

²University of Oklahoma, Norman, Oklahoma 73019

³Purdue University, West Lafayette, Indiana 47907

⁴University of Rochester, Rochester, New York 14627

⁵Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309

⁶Southern Methodist University, Dallas, Texas 75275

⁷Syracuse University, Syracuse, New York 13244

⁸Vanderbilt University, Nashville, Tennessee 37235

⁹Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

¹⁰Wayne State University, Detroit, Michigan 48202

¹¹California Institute of Technology, Pasadena, California 91125

¹²University of California, San Diego, La Jolla, California 92093

- ¹³University of California, Santa Barbara, California 93106
¹⁴University of Colorado, Boulder, Colorado 80309-0390
¹⁵Cornell University, Ithaca, New York 14853
¹⁶University of Florida, Gainesville, Florida 32611
¹⁷Harvard University, Cambridge, Massachusetts 02138
¹⁸University of Hawaii at Manoa, Honolulu, Hawaii 96822
¹⁹University of Illinois, Urbana-Champaign, Illinois 61801
²⁰Carleton University, Ottawa, Ontario, Canada K1S 5B6
and the Institute of Particle Physics, Canada
²¹McGill University, Montréal, Québec, Canada H3A 2T8
and the Institute of Particle Physics, Canada
²²Ithaca College, Ithaca, New York 14850
²³University of Kansas, Lawrence, Kansas 66045
²⁴University of Minnesota, Minneapolis, Minnesota 55455
²⁵State University of New York at Albany, Albany, New York 12222

Although several modes of radiative and hadronic $\Upsilon(1S)$ decays with multiparticle final states have previously been observed, no radiative decays of the $\Upsilon(1S)$ into a photon and two hadrons have yet been reported. Such final states have provided the most direct evidence for two-body radiative J/ψ decays, which are well established [1] at the 10^{-3} level. To extrapolate these to the Υ , the charge coupling to the photon and the mass of the quark propagator predict a suppression of order $[(q_b/q_c)(m_c/m_b)]^2 \sim 1/40$. More sophisticated calculations can be found in the literature [2].

The radiative decays of the $\Upsilon(1S)$ can provide information on exotic states, including WIMP's and axions [3,4]. The observation of resonances in the two-body invariant mass spectrum opposite photons is also one of the ways of establishing possible glueball candidates in radiative quarkonium decays, using the fact that the emission of the photon leaves a recoiling “glue rich” environment from which to form them. Radiative decays of the $\Upsilon(1S)$ with charged final state hadrons have been studied by many experimental groups, including ARGUS [5], CLEO [6], and MD-1 [7]. In the CLEO analysis, the decay modes $\Upsilon(1S) \rightarrow \gamma X$; $X \rightarrow \pi^+\pi^-$, K^+K^- and $p\bar{p}$ were investigated. As noted in that ref. [6], the only region of the dipion invariant mass distribution suggestive of an excess above background was in the interval 1.2-1.6 GeV, where ten signal events were counted; the scaled background in the same region corresponded to two events. In this Letter, we extend the previous CLEO analysis, using a new data set and exploiting many improvements in the performance of the CLEO II detector [8], which operates at Cornell Electron Storage Ring (CESR). We also present first results for the all-neutral final state $\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0$.

The signal data used in this analysis were collected on the $\Upsilon(1S)$ resonance at a center-of-mass energy $E_{cm} = 9.46$ GeV, corresponding to an integrated luminosity of 78.9 pb^{-1} . Data taken at $E_{cm} \cong 10.52$ GeV, just below the $\Upsilon(4S)$ resonance, were used to subtract the $e^+e^- \rightarrow \gamma X$ events due to non- $\Upsilon(1S)$ production under the resonant $\Upsilon(1S)$ peak; this sample corresponds to an integrated luminosity of 500.4 pb^{-1} . We search for events in both our $\Upsilon(1S)$ (signal) and continuum (background) datasets compatible with the kinematics for the process $\Upsilon(1S) \rightarrow \gamma\pi\pi$. Separate criteria are applied for the cases $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ and $\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0$.

Candidate events for the $\gamma\pi^+\pi^-$ final state are selected as follows. There must be exactly two oppositely-charged tracks observed in the detector. If the ratio of a track's energy deposited in the calorimeter to its momentum measured in the drift chambers is greater than 0.85, the track is identified as an electron and the event vetoed from further consideration. At least one of the charged tracks must satisfy the kinematic requirements for muon identification, defined in terms of a track's polar angle (θ) and its momentum (p) as $|\cos\theta| < 0.7$ and $p > 1.0 \text{ GeV}/c$. Any track satisfying these criteria and also producing associated hits in the muon chambers is identified as a muon, and the event is similarly vetoed. There must be exactly one electromagnetic shower in the good barrel region of the calorimeter ($|\cos\theta| < 0.71$) with energy exceeding $0.4 \times E_{cm}$. This shower must have an energy deposition profile consistent with that of a photon, and also not match, within 15 degrees, the position of any charged track extrapolated into the calorimeter. Additional showers, presumed to be either noise or split-offs from the tracks propagating into the detection volume of the calorimeter, are allowed provided their measured energies are each less than 500 MeV. The sum of the energy of the highest energy photon candidate plus the energies of the drift chamber tracks must, under the $\pi^+\pi^-$ hypothesis, lie within three standard deviations in energy resolution (σ_E) of E_{cm} . Typically, we find $\sigma_E \cong 80 \text{ MeV}$. $\Upsilon(1S) \rightarrow \gamma K^+K^-$ and $\Upsilon(1S) \rightarrow \gamma p\bar{p}$ events, although not yet observed, are, if misinterpreted, more likely to fail this energy-conservation requirement than true $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ events. The magnitude of the net momentum vector of the event must be within three standard deviations (σ_p) of zero; σ_p takes into account the resolutions on the two tracks and

the high energy photon and is typically 80 MeV/c. We require that the opening angle ϕ between the two charged tracks satisfy the condition $\cos \phi > -0.95$.

The momentum of each charged track recoiling against the high energy photon is typically ≈ 2 GeV/c, beyond the momentum range for which the CLEO detector can cleanly separate pions from kaons or protons. We therefore only require that the available dE/dx particle identification information be consistent with the pion hypothesis. We have, nevertheless, performed dedicated searches for $\Upsilon(1S) \rightarrow \gamma K^+ K^-$ and $\Upsilon(1S) \rightarrow \gamma p \bar{p}$. In neither case was any signal above background observed.

Candidate $\gamma\pi^0\pi^0$ events must have no charged tracks. The requirements on the high-energy photon in the event are identical to the case of $\gamma\pi^+\pi^-$. Neutral pions are defined as combinations of two showers in the electromagnetic calorimeter with an invariant mass within five standard deviations of the nominal π^0 mass. All photons in π^0 reconstruction must also satisfy a minimum energy requirement ($E_\gamma > 50$ MeV), and have an energy deposition pattern consistent with true photons. The four-momentum conservation requirements are also identical to the charged pion case.

We use a GEANT-based [9] detector simulation to determine the efficiency for reconstructing a radiative $\Upsilon(1S)$ event, as a function of dipion mass, for each final state studied. Candidate $\Upsilon(1S) \rightarrow \gamma X$ events are generated by LUND Monte Carlo [10] with flat distribution over the entire kinematically allowed m_X regime. The recoil system decays isotropically, and the final state pions are propagated through the detector. The overall event selection efficiency (ϵ) for the $\gamma\pi^+\pi^-$ final state varies smoothly from $\epsilon \cong 33\%$ at threshold ($m_{\pi^+\pi^-} = 2m_\pi$) to a maximum of $\epsilon \cong 41\%$ at $m_{\pi^+\pi^-} \cong 2$ GeV. By comparison, the event reconstruction efficiency for the $\gamma\pi^0\pi^0$ final state grows rapidly from zero at threshold to $\epsilon \cong 30\%$ at $m_{\pi^0\pi^0} \cong 1$ GeV and then smoothly falls to $\epsilon \cong 28\%$ at $m_{\pi^0\pi^0} \cong 2$ GeV.

The invariant mass of the recoiling hadrons for candidate events is presented in Fig. 1(a) (charged pions) and Fig. 1(b) (neutral pions), for both the $\Upsilon(1S)$ resonance data and the continuum data. The continuum data have been properly scaled to the $\Upsilon(1S)$ data, taking into account the differences in the luminosity of our signal and background event samples, the expected $1/E_{bm}^2$ energy dependence of the QED cross section, and the relative event selection efficiency for the $\Upsilon(1S)$ and the continuum data.

Prominent in Fig. 1(a) is a large ρ^0 signal as verified experimentally [5–7] by other analyses. Backgrounds due to $e^+e^- \rightarrow \gamma\rho^0$, $\rho^0 \rightarrow \pi^+\pi^-$ are expected to dominate the $\gamma\pi^+\pi^-$ analysis. Comparing the acceptance and luminosity-corrected signals observed in both the $\Upsilon(1S)$ and continuum data, we note that the $\gamma\rho^0$ rate observed on the $\Upsilon(1S)$ is consistent with the yield from the continuum data. Subtracting the scaled continuum dipion mass distribution from the resonant $\Upsilon(1S)$ mass distribution in the “ ρ -rich region” ($m_{\pi\pi} < 1.0$ GeV), we obtain 1.4 ± 21.0 events, consistent with zero. Also evident is a small enhancement at $m_{\pi\pi} \approx 400$ MeV which is largely from misidentified $e^+e^- \rightarrow \gamma\phi$; $\phi \rightarrow K^+K^-$ events. The scaled continuum data are consistent with the resonant $\Upsilon(1S)$ yields.

We note a significant excess of events in the $\Upsilon(1S)$ data sample over background in both the $\gamma\pi^+\pi^-$ and the $\gamma\pi^0\pi^0$ final states. Performing a bin-by-bin continuum subtraction, we obtain an excess of 47.0 ± 9.3 (9.0 ± 3.0) events for the $\pi^+\pi^-$ ($\pi^0\pi^0$) data, integrated over $m_{\pi\pi} \geq 1$ GeV. We attribute these excesses to the decays $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ and $\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0$, respectively. Based on the total number of $\Upsilon(1S)$ events in our sample (1.86×10^6), and correcting for the efficiencies as a function of invariant mass, we obtain $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-) = (6.3 \pm 1.2 \pm 1.3) \times 10^{-5}$ and $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma\pi^0\pi^0) = (1.7 \pm 0.6 \pm 0.3) \times 10^{-5}$, for $m_{\pi\pi} \geq 1$ GeV, in which the second error is systematic (to be




FIG. 1. Dipion invariant mass for the $\Upsilon(1S)$ data, with scaled continuum data (shaded) overlaid, for the final states $\gamma\pi^+\pi^-$ (a) and $\gamma\pi^0\pi^0$ (b).

described later).

Whereas the statistics in the background-subtracted $m_{\pi^0\pi^0}$ mass distribution are too poor to show any obvious structure, the excess in the charged dipion mode is apparent in the region $m_{\pi^+\pi^-} \approx 1.0$ -1.4 GeV. The most prominent resonance in this mass range observed in radiative J/ψ decays is the $f_2(1270)$ [1], for which $\mathcal{B}(J/\psi \rightarrow \gamma f_2(1270)) = (1.38 \pm 0.14) \times 10^{-3}$. If we assume that the excess in this interval is due to $\Upsilon(1S) \rightarrow \gamma f_2(1270)$, and neglecting any possible interference effects with other processes producing the $\gamma\pi^+\pi^-$ final state, we can perform a fit to the background-subtracted on-resonance dipion invariant mass spectrum and thereby determine the possible level of $\Upsilon(1S) \rightarrow \gamma f_2(1270)$, as shown in Fig. 2. In performing this fit, we use a spin-2 Breit-Wigner signal function with the mean and width parameters fixed to the established [1] $f_2(1270)$ values, using the interval 0.6–1.8 GeV. Such a fit yields 34.8 ± 9.7 events with a χ^2 per degree of freedom of 12.8/23. If, instead, we allow the mass and width to float, we obtain a yield of $30.1^{+9.9}_{-9.3}$ events, with a fitted mass of (1.28 ± 0.02) GeV and a width of (100^{+80}_{-40}) MeV. Assuming no other contributions to the spectrum in Fig. 2, the corresponding efficiency-corrected product of branching fractions would be $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma f_2(1270)) \times \mathcal{B}(f_2(1270) \rightarrow \pi^+\pi^-) = (4.6 \pm 1.3^{+1.6}_{-1.5}) \times 10^{-5}$, which gives [1] $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma f_2(1270)) = (8.1 \pm 2.3^{+2.9}_{-2.7}) \times 10^{-5}$. The likelihood that the excess in this region is due to an upward fluctuation of background is determined to be less than 0.01%.

If this excess is due to $\Upsilon(1S) \rightarrow \gamma f_2(1270)$, then, by isospin, we expect to also observe $\Upsilon(1S) \rightarrow \gamma f_2(1270)$, $f_2(1270) \rightarrow \pi^0\pi^0$ at half the charged rate. When compared to the fitted result in the charged dipion case, the excess of 10.8 ± 3.3 events observed is consistent, after efficiency-correction, with this expectation. The probability of the continuum background fluctuating up to the $\pi^0\pi^0$ signal is negligible. Non- f_2 contributions, if any, are difficult to assess with the limited statistics of the signal sample.

Although there is no resonance with $m \approx 1.05$ GeV expected in our sample, we nevertheless note an apparent enhancement in this mass region in Fig. 2. To investigate this further, we have allowed

for a second Breit-Wigner in our fit, with the values of mass and width for this second Breit-Wigner allowed to float, but with the f_2 parameters again constrained to the established values (Fig. 2). We then obtain 20.5 ± 12.3 events for this second Breit-Wigner, at a mass of (1.05 ± 0.02) GeV and a fitted width of (100 ± 90) MeV; the putative $f_2(1270)$ yield correspondingly drops to 29.7 ± 11.0 events. The overall χ^2 per degree of freedom in this second fit improves to 6.2/20. We note that although the level of the excess at 1.05 GeV is at the 1.7 standard deviation level, the likelihood that the excess in the 1.28 GeV mass region is an upward fluctuation of background is still small (less than 0.1%).




FIG. 2. Fit of the continuum-subtracted $\Upsilon(1S)$ dipion invariant mass spectrum to a single $f_2(1270)$ only (dashed line, and blown up in inset) with overlay of fit to $f_2(1270)$ plus a possible second resonance at $m_{\pi^+\pi^-} \approx 1.05$ GeV (shaded region under the solid line). Note that the $f_2(1270)$ yield is relatively insensitive to the addition of the second resonance.

We have considered possible contamination to our signal from the process $\Upsilon(1S) \rightarrow \gamma\mu^+\mu^-$. This is evaluated by selecting, rather than vetoing, events having a high energy photon and two charged tracks, in which there are hits in the muon chambers matched to at least one of the charged tracks. For such a search, the continuum-subtracted $\Upsilon(1S)$ data yields 18 ± 19 $\Upsilon(1S) \rightarrow \gamma\mu^+\mu^-$ event candidates. Knowing that the maximal inefficiency for $\gamma\mu^+\mu^-$ events is 30%, we determine that the contribution from the $\gamma\mu^+\mu^-$ final state to our signal $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ sample has a central value less than 5.4 events, and consistent with zero; we include this in our systematic uncertainty.

The decay $\Upsilon(1S) \rightarrow \pi^0\pi^+\pi^-$, although not yet observed, could produce background to the $\gamma\pi^+\pi^-$ final state in the cases in which the π^0 decays either very asymmetrically (resulting in one very high energy shower with energy almost equal to the energy of the parent π^0), or produces a π^0 in which both decay photons are approximately collinear, and cannot be distinguished. In the latter case, the photon showers overlap and merge into a single detected calorimeter shower. We have conducted a dedicated search for the decay $\Upsilon(1S) \rightarrow \pi^0\pi^+\pi^-$, with identical requirements on the π^0 candidate, on the tracks of the two charged pion candidates, and on four-momentum conservations as in the $\gamma\pi\pi$ analysis. This search was consistent with zero events being found. The associated

upper limit is $\mathcal{B}(\Upsilon(1S) \rightarrow \pi^0 \pi^+ \pi^-) < 1.84 \times 10^{-5}$ at 90% confidence level. Based on this null result and the probability to misinterpret a “merged” π^0 as a photon, we would expect fewer than 3.4 events contamination from the decay $\Upsilon(1S) \rightarrow \pi^0 \pi^+ \pi^-$ in our $\gamma \pi^+ \pi^-$ event sample over the entire kinematically allowed dipion invariant mass range. The net contribution from $\Upsilon(1S) \rightarrow \pi^0 \pi^+ \pi^-$ events in which the π^0 decays asymmetrically is determined to be less than 0.7 events.

We note that two pions produced in $\Upsilon(1S) \rightarrow \gamma f_2(1270)$, $f_2(1270) \rightarrow \pi^+ \pi^-$ will have a characteristic angular distribution, due to the tensor nature of the $f_2(1270)$. We have correspondingly fit the helicity angle distribution (defined as the angle between one of the pions and the dipion parent measured in the dipion rest frame) for the mass interval 1.2-1.4 GeV, after subtracting out the contribution from the $\gamma \rho$ final state. Such a fit gives confidence levels of 48%, 35%, 0.0%, and 0.1% for the system recoiling against the photon to be a tensor, scalar, vector or axial vector, respectively. Although inconclusive on its own, this spin-parity analysis of the dipion system strongly favors a tensor or scalar assignment for the dipion system, and rules out a vector or axial vector interpretation.

For the measurement $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$, systematic uncertainties are due primarily to the muon veto used to suppress the $\gamma \mu^+ \mu^-$ final state (12% relative error), uncertainties in our total efficiency (5%, arising mainly from event triggering uncertainties), and our uncertainty in the total number of $\Upsilon(1S)$ events (3%). Because we have assumed that the photon angular distribution is isotropic in our Monte Carlo event generator, there is an additional uncertainty (16%) from our extrapolation to the region $|\cos \theta_\gamma| > 0.71$. For the possible $\Upsilon(1S) \rightarrow \gamma f_2(1270)$ signal, we have an additional systematic error (20%) due to the fitting procedure used to extract the signal, including the possible effect of the apparent enhancement in the region $m_{\pi^+ \pi^-} \approx 1.05$ GeV, and asymmetric uncertainties due to the possible interference between events from the $\Upsilon(1S) \rightarrow \gamma f_2(1270)$ excess and $\pi^+ \pi^-$ pairs not associated with either the ρ^0 or the resonant enhancement ($^{+20}_{-15}$)%.

For the $\gamma \pi^0 \pi^0$ final state, primary uncertainties in our integrated measurement $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0)$ are due to the possible anisotropy of $\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$ decay (16%), π^0 finding (8%), trigger efficiency (4%), and the number of $\Upsilon(1S)$ events (3%).

In summary, we have made the first observation of the radiative decay $\Upsilon(1S) \rightarrow \gamma \pi \pi$ in both charged and neutral modes. Restricted to $m_{\pi\pi} \geq 1.0$ GeV, we obtain $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-) = (6.3 \pm 1.2 \pm 1.3) \times 10^{-5}$, and $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0) = (1.7 \pm 0.6 \pm 0.3) \times 10^{-5}$.

The $\pi^+ \pi^-$ mass and helicity angle distributions are suggestive of $f_2(1270)$ production as a source. Under the $\Upsilon(1S) \rightarrow \gamma f_2(1270)$ assumption, the efficiency-corrected product of branching fractions of this enhancement corresponds to $\mathcal{B}(\Upsilon(1S) \rightarrow \gamma f_2(1270)) \times \mathcal{B}(f_2(1270) \rightarrow \pi^+ \pi^-) = (4.6 \pm 1.3^{+1.6}_{-1.5}) \times 10^{-5}$. In the $\pi^0 \pi^0$ mode, the net yield relative to the charged mode and the shape of the $\pi^0 \pi^0$ mass spectrum are also consistent with $\Upsilon(1S) \rightarrow \gamma f_2(1270)$. This is approximately twice the rate that would be expected by the $[(q_b/q_c)(m_c/m_b)]^2$ scaling from $\mathcal{B}(J/\psi \rightarrow \gamma f_2(1270))$.

Acknowledgments

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the National Science Foundation, the U.S. Department of Energy, Research Corporation, the Natural Sciences and Engineering Research Council of Canada, the A.P. Sloan Foundation, the Swiss National Science Foundation, and the Alexander von Humboldt Stiftung.

REFERENCES

- [1] R.M. Barnett *et al.*, Particle Data Group, Phys. Rev. **D 54**, 41 (1996).
- [2] J.G.Körner *et al.*, Nucl. Phys. **B 229**, 115 (1983).
- [3] P.Fayet and J.Kaplan, Phys. Lett. **B 269**, 213 (1991).
- [4] CLEO Collaboration, R. Balest *et al.*, Phys. Rev. **D 51**, 2053 (1995).
- [5] ARGUS Collaboration, H. Albrecht *et al.*, Zeit. Fur. Phys. **C 42**, 349 (1989).
- [6] CLEO Collaboration, R.Fulton *et al.*, Phys. Rev. **D 41**, 1401 (1990).
- [7] MD-1 Collaboration, S. Baru *et al.*, Phys. Rep. **267**, 71 (1996).
- [8] CLEO Collaboration, Y. Kubota *et al.*, Nucl. Inst. Meth. **320**, 66 (1992).
- [9] R. Brun *et. al.*, GEANT v. 3.15, CERN Report No. CERN CC/EE/84-1 (1987).
- [10] T. Sjostrand and M. Bengtsson, JETSET v. 7.3, Comput. Phys. Commun. **43**, 367 (1987).